

УДК 796. 01: 159.9; 612.821-053.6

Psycho-Physiological Types of Adaptation in the Assessment of Professional and Sports Selection

Olga N. Moskovchenko*

Siberian Federal University

79 Svobodny, Krasnoyarsk, 660041 Russia ¹

Received 2.09.2011, received in revised form 9.09.2011, accepted 16.09.2011

Preliminary research allowed to determine the types of psycho-physiological adaptation on the basis of analysing the non-parameter algorithms of the automatic classification (Lapko, 1993); four types of psycho-physiological adaptation have been identified.

It is expected to elaborate previously determined basic criteria which integrate neuro-dynamic profiles of adaptation characterising the individual peculiarities of the central nervous system, its power and lability, indicators of the heart rate variability. As additional criteria we offer differentiated morphofunctional types of physical development and typological types of personality determined in accordance with Hans Eysenk's Personality Chart (1972). Such an approach will give an opportunity to consider the types of psychophysical adaptation in the unity of personal features and genetically natural potential providing individual and psychological peculiarities of the personality.

A brief characteristics of each type of psycho-physiological adaptation was given, thus contributing to understanding of the personality individual features and providing the opportunity to forecast behaviour reactions.

The studies have enhanced understanding of the types of psycho-physiological adaptation, determination of the forecasting criteria which allow to predict the behaviour in extreme situations, in terms of educational, industrial and sports activities. The types of psycho-physiological adaptation can be considered when assessing the socio-psychological adaptation of students, as well as the criteria for professional and sports selection.

Keywords: types of psycho-physiological adaptation, sprinter, stayer, mixed type, unsteady type, selection criteria.

Introduction

In modern world psychology the main part of research studies is devoted to the issues of psycho-physiological adaptation and is aimed at studying the psychological adaptation mechanisms in the conditions of acclimatisation. In psychological studies the basic characteristics of interpersonal

relationships, their types and mechanisms of appearance are determined. (Bodrov, 1995; Kolyshkin, 1997; Baker, 1969; Budd, 1973; Wallhoffer, 1976 et al.).

In recent years some authors (Efimova, 2003; Tiunova, 2006, Rodionov, 2004 et al.) consider psycho-physiological criteria as the basis for

* Corresponding author E-mail address: moskovchenko7@mail.ru

¹ © Siberian Federal University. All rights reserved

students' health and also as the correction of unfavourable psycho-emotional states in physical education and sports.

However, it should be noted that at present neuro-physiological peculiarities of human individual adaptation are poorly known (Bekhtereva, 1974; Nebylitsyn, 1978; Soroko, 1978). In some publications different psycho-physiological types are determined by formal features (age, gender, interpersonal relationships, etc.).

“Psycho-physiological portraits” proposed by V.L. Bakaleynikova (1986) are defined by the functional features of the psychomotor activity, thus being more informative, but not solving the problem of the psycho-physiological adaptation in general.

Researchers have not been paying significant attention to the issues of classifying the types of adaptation. For the first time the ecological types of adaptation were identified by V.P. Kaznacheyev and S.V. Kaznacheyev in 1986. In other papers concerning the adaptation types the researchers refer to the authors mentioned above.

Nevertheless, in our opinion the psycho-physiological adaptation (PPA) should be founded not only on the basic mental functions, but on the physiological factors, compensative adaptive mechanisms of autonomic regulations of the heart rate and the central nervous system taking into consideration typological peculiarities of the personality, thus providing functional reserve of the individual's body. Such an approach coincides with the opinion of F.B. Berezin (1988).

Therefore, the types of the human psycho-physiological adaptation should be determined not only by the formed psycho-physiological features of the personality and parameters of autonomic homeostasis, but also by the state of the body physiological systems characterised by neuro-dynamic profiles determined genetically and fixed by the phenomic conditionality which

provides mental and motor activity, types of individual's adaptive behaviour and has practical implication.

During our studies of the psycho-physiological adaptation in the process of educational activities and sportsmen training we identified contradictions between determining the types of adaptation and the existing variety of opinions regarding this issue. Absence of unambiguous opinions regarding the criteria of the PPA types has determined the direction of our further research in this field. As the basis for this work we took the validation of the PPA types from the point of view of the integrating role of the specific functional systems which not only provide adaptive reactions by interacting with each other, but also coordinate the activities of psycho-physiological functions.

Methods of examination and conditions of the experiment

In the examination 12,000 first-year students of basic medical groups from different universities of Krasnoyarsk took part on the basis of the interuniversity valeological centre (KSTU – SibFU), as well as skilled sportsmen of different kinds of sports, among them 5 honoured masters of sports, 12 masters of sports of the international level, 53 masters of sports and 25 candidates for masters of sports. Physiological and psycho-physiological functions were studied in the laboratory and in the training camps. The morning examination was done from 9.00 to 12.00, the evening examination – from 16.00 to 18.00. One procedure of the examination lasted for 3 days, the diagnostics was carried out according to the stages of the developed pattern.

At the first stage the indicators characterising neuro-dynamic profiles were measured: omegametry (DLOP – dominant levels of omega-potential); methods for evaluating the functional

state of the central nervous system: CFLF-critical frequency of light flashes, BEP – bio-energy potential, an analogue of galvanic skin response, MFM – the maximum frequency of movements (tapping test); VMR – visual-motor reaction; Lüscher test using the author's expert system "LUSY". The certificate of registration of computer programmes No. 2002610463. Moscow, 2002.

At the second stage the indicators of the evaluation of autonomic regulation of the heart rate mechanisms were recorded using the hardware-software complex (Aldonin, 1998), Questionnaire of H. Eysenck.

At the third stage in order to determine the morphofunctional types of physical development (MFTPD) the anthropometric parameters (height, weight, lung capacity, hand dynamometry and circumference of the chest) were measured. The massiveness of the skeleton was calculated by the largest circumference of the chest and in accordance with the calculation formulas.

Complex assessment of the PPA was done using the author's expert system "KOMPEA". The certificate of computer programmes' registration in the Specific Fund of Algorithms and Programmes of the Ministry of Defence of the Russian Federation No. ОФАП – 1824 No. r.p.50200200003, Moscow, 2002.

For students the measurements were made during 4 semesters (the 1st -2nd year students) to determine the duration of the adaptation periods. For sportsmen the measurements were done at the beginning, in the middle and at the end of the training process.

Statistical processing of the experimental data and mathematical modelling were done using the dialog package of "NPCL" programme developed by G.S. Vysotskaya (1998) at the Information and Computing Centre of the Siberian Branch of the Russian Academy of Sciences and "STATGRAF" package.

Results and discussion

Our preliminary research allowed to determine the typical features of the psycho-physiological adaptation types (Table 1).

At the evaluation of psycho-physiological mechanisms of adaptation the determining criteria were: psychomotor function and autonomic homeostasis (Moskovchenko, 1999, 2000, 2004). Moreover, each of the types differed from each other by the quantitative criteria of the studied features. Two types were designated in accordance with the classification of V.P. Kaznachejev (1986) as a "sprinter" and a "stayer", the other two were classified by the author. The intermediate type was classified as a "mixed" type, and the type with unstable neuro-dynamic profile as an "unsteady" type.

Our further research relied on the theory of functional systems (Anokhin, 1980; Sudakov, 1983); on the structural and functional organisation of the brain systems in ensuring the mental and motor activity of the human within the framework of the theory (Aladzhlova, 1979, Bekhtereva, 1974), as well as on the role of ultraslow physiological processes of the brain (Sychev, 1980; Ilyukhin, 1982; Moskovchenko, 1997, 2004). We also considered the methodology offered by V.P. Kaznachejev (1980, 1986) regarding the mutual influence of psycho – and somatotype, which will allow to reveal not only different behavioural responses in the structure of the adaptive mutations, but also health hidden reserves.

Basing on the mathematical modelling with implementation of the nonparametric algorithms of automatic classification, for the first time it became possible to identify a previously unknown pattern in the selection of quantitative criteria, six neuro-dynamic profiles and five morphofunctional types of physical development which differ from each other by the reproduction of the generated adaptive programmes.

Table 1. Typical features of the psycho-physiological adaptation types

Types of the psycho-physiological adaptation	Criteria of assessment of power and lability of the nervous system, psycho-motor skills and autonomic coefficient by Lüscher X, \pm d						Indicators and oscillations of the variation pulsemetry		
	CFLF, Hz	BEP, relative unit.	MFM 1 strike/10s	VMR, ms	Weight category by cytotoxic lymphocytes, units	$\Delta X, c$	AM ₀ /Δx	Tension index	
Sprinter	48-60 \pm 1.02-1.34	1-1.4 \pm 0.002-0.01	75-90 \pm 3.32-4.34	150-200 \pm 5.65 -6.21	1-1.6 \pm 0.02-1.2	0.23-0.45 0.20-0.30 \pm 0.02-1.33 \pm 0.02-1.21	120-140 100-115 \pm 5.89-6.76 \pm 4.25-5.8	60-90 \pm 2.18-2.54	
Stayer	35-48 \pm 1.52-1.77	0.85-1 \pm 0.54 -0.98	50-65 \pm 1.71-1.65	200-250 \pm 7.26-8.56	0.95-0.80 \pm 0.01-1.02	0.25-0.40 0.40-0.45 \pm 2.32-2.89 \pm 2.43-1.94	120-140 \pm 3.45-4.81	100-120 \pm 1.67-2.51	
Mixed type	40-50 \pm 1.27-1.44	1.0-0.85 \pm 0.12-0.58	70-85 \pm 2.02-1.34	180-220 \pm 1.03-2.62	1-2.0 0.8-0.99 \pm 0.02-0.62 \pm 0.01-0.72	0.20-0.38 0.30-0.40 \pm 0.01-0.2 \pm 0.02-0.46	100-120 125-140 \pm 2.66-3.48 \pm 4.02-5.4	60-130 \pm 1.09-4.26	
Unsteady type	32-37 \pm 1.52-1.58	0.84-0.50 \pm 0.02-0.37	40-59 \pm 1.89-1.94	250-345 \pm 13.42-18.76	2.1-3.0 0.7-0.8 \pm 1.65-1.80 \pm 1.33-1.34	0.17-0.29 \pm 0.01-0.02	140-180 \pm 2.59-3.71	120-190 \pm 3.48-5.35	

Table 2. Types of psycho-physiological adaptation

Types of psycho-physiological adaptation	Features of the typological peculiarities, extraversion-introversion scale	Neuro-dynamic profiles of adaptation	Morphofunctional type of physical development	Functional state of the CNS		
				Lability	Mobility	Level of energetic mobilization
Sprinter	Extravert, choleric, sanguine	Steady, hyper steady	II, III	Very high	Very high	High or good
Stayer	Introvert, phlegmatic, melancholic	Steady, hyper steady, hypo steady	III, IV, I	Good	Good	High or good
Mixed type	Extravert, introvert	Steady, hyper steady, hypo steady	II, III, IV	Very high	High	Very high
Unsteady type	Neurotic character	Unsteady, hypo-unsteady, hyper-unsteady	I, IV, V	Medium or low	Low	Medium or low

The morphofunctional types of the physical development (MFTPD) have been differentiated as follows: I type – disharmonious, with the asthenic body type; II type – harmonious, with the normosthenic body type; III type – harmonious, with the muscular body type; IV type occupies an intermediate position between II and V types, and have not been associated with any classical notion characterising any body type; V type – disharmonious, with the hypersthenic body type.

High reliability between MFTPD ($p < 0.001$), its variability depending on the place of residence for one and the same socio-demographic group has been determined, thus forming the basis for consideration of the morphofunctional types as the component of the morphofunctional properties of the body determining the adaptive functions of adaptive capacity.

According to some authors (Kaznacheyev, 1986; Kuraev, 2002; Moskovchenko, 2004, et al.) the morphofunctional parameters are a genetic factor and are fixed by phenotypic conditionality providing mental and motor activity at the individual level, which serve as the basis for

considering MFTPD in assessing the PPA types (Table 2).

Such interdependence characterises general biological significance of adaptive functional reserves of the body, provides various adaptive reactions to environmental conditions, industrial, educational and sports activities, reflects behavioural reaction to any kind of influences.

To the “sprinter” type persons with II and III MFTPD are referred, with the steady or hyper steady neuro-dynamic profile. The normosystolic tone – eutonia (an indicator of the balance of the autonomic nervous system, heart rate variability) was observed among 80 % – 20 %. Parasympathetic and moderately sympathetic influences predominate indicating the reserve capacity of the body and corresponding with the data of R. Baevsky (1986). The “sprinter” type is mostly typical for extroverts-leaders with high motivation, as in the activity itself, so in the choice of the profession.

Fairly high lability and mobility of the nervous processes contribute to the rapid switching from one type of activity to another one; balance and

strength of the nervous processes allow to get quickly mobilised for work of any type in normal and extreme conditions. The “sprinters” easily withstand short-term stressful effects and are able to quickly switch for new adaptive programmes.

For the “sprinter” the features of sanguine and choleric temperament are typical; both types are characterised by a strong nervous system but with its various mobility or lability, various degrees of energy mobilisation. Long-term dynamics has demonstrated that the “sprinters” with II and III MFTPD, the sanguine temperament and the steady neuro-dynamic profile are more adapted to the ecological, climatic and zone conditions, stressful situations. Sportsmen are able to maintain difficult-to-coordinate movements during competitions, even if they are tired. Such qualities as persistence, energy, innovation, competitive spirit, emotional stability, assertiveness, cooperation, capability to show realism in their claims dominate in the structure of the motivation for the sports activity.

The “sprinter” is characterised by high levels of social and psycho-physiological adaptation which allows to meet the requirements of the higher educational institution (for students) successfully, contributes to sports activity and self-realisation of the person.

To the “stayer” type the persons with a stable, hypo- or hyper steady neuro-dynamic profile refer. High correlation has been noted for the persons with a phlegmatic temperament with IV MFTPD ($r = 0.72$) and for the persons with a melancholic temperament with I ($r = 0.86$), insignificant correlation respectively with II ($r = 0.46$) and IV ($r = 0.38$) MFTPD. In the stayers’ group the autonomic balance – eutonia has been observed in 43 % of cases, among phlegmatic persons sympathicotony prevailed in 37 % of cases, while among melancholic persons in 20 % an increased parasympathetic influence – vagotonia was noted.

The “stayer” type is characterised by relatively low or high values of the dominant level of the omega potential (DLOP). At low values (12-19 mV) it is a hypo steady type with good adaptation in favourable social conditions, in extreme situations adaptive responses were reduced, psycho-physiological processes became unsteady resulting in disorganization of behaviour, unnecessary anxiety appeared with and without a reason. At high values of DLOP (44-48 mV) it is a hyper steady type. Individual adaptation depended on the mental states, motivation and conditions of the social environment. Often emotional reaction to educational or other activities was higher due to overestimation of own capabilities. During training it is necessary to use not only the method of the story, but also the method of the show. In extreme conditions high probability of failure of adaptive mechanisms appears that can lead to disorganization of behaviour, especially during decision making.

The “stayer” with the average DLOP (20 – 40 mV) is characterised by high mobility of nervous processes, but by high lability, or vice versa. The “stayers” are capable of performing various actions for a long period of time and in any situation, but at a slower pace if compared with the “sprinters”. They endured quite good short-term stressful effects at positive socially significant conditions (constant support, approval, advice). Socio-psychological adaptation in extreme conditions is impeded, prolonged exposure to inadequate conditions led to the breakdown of adaptation. As a rule, the “stayers” were active in the personal life. The “stayer” with a melancholic temperament is characterised by social passivity, he tries to avoid responsibility seeking protection from unnecessary conflicts.

To the “mixed” type the persons with a choleric, sanguine and phlegmatic character are referred, with a stable, hypo- or hyper steady neuro-dynamic profile, but with a tendency to

periodic disturbance of homeostatic systems (instability of the omega gram maintenance in the amplitude-time structure from day by day). A significant proportion – 75 % consisted of the eutonic persons, among 25 % the dominant influence of the sympathetic or parasympathetic link in the regulation of the sinus heart rate was observed. II, III and IV MFTPD indicated the harmony of the morphotype. The “mixed type” with a sanguine temperament is characterised by good mobility and high lability of the nervous processes, or vice versa. They were able to perform a variety of actions equally well in all conditions, as in the fast, so in the slow pace. High emotional stability allowed to adapt to prolonged exposure to extreme factors in unfavourable conditions.

The “mixed type” with a choleric temperament is characterised by good mobility of the nervous processes, they are highly active, impulsive, but the elements of incoherence were present in activities, they could not concentrate on the essential part. Excessive impulsivity in prolonged unfavourable conditions often led to mistakes, disorganization of behaviour, overstrain of adaptive mechanisms. Persons with a phlegmatic temperament endured the stressing effects quite well at the positive socially significant conditions. If these conditions were violated the psycho-physiological adaptation decreased. Therefore, one can conclude that for the “mixed” type the social status plays an important role in the process of psycho-physiological adaptation.

The “unsteady type” includes all those who had low (8-16 mV) or high (42-48mV) DLOP values, with stable violation of chemometabolic processes (transition of ω -potential from the initial level to the value up to 25 % after functional loading) indicating unsteady adaptive capacities of the body at physical or mental workload, which displayed the reduction of mental and

physical functions of the body reserves in extreme conditions and was accompanied by a decrease of efficiency, intolerance to long-term exercise, instability and insecurity in education or sports activities. Significant physical activities led to rapid depletion of the central regulatory mechanisms and disruption of adaptation. For the persons of the “unsteady type” significant correlation is marked with I ($r = 0.82$), IV ($r = 0.79$) and V ($r = 0.66$) MFTPD at ($p < 0.05$) and, respectively, insignificant correlation with II ($r = 0.26$) and III ($r = 0.38$) type at ($p < 0.005$). We judged about the degree of adaptation mechanisms strain by the expressed changes in the autonomic balance of sympathetic and parasympathetic influence.

The “unsteady type” is characterised by medium or low lability of the nervous processes and the middle or the low level of energy mobilisation. Individuals with low values of DLOP were able to perform long-term repetitive actions at the medium pace and in the familiar environment. In extreme situations (a test or an exam) the concentration of attention and speed of information processing were maintained for a short period of time, then a tendency to quick tiredness appeared, which led to errors. Among persons with a high value of DLOP social and psycho-physiological adaptation was impeded due to manifestations of slow response and increased anxiety, excessive suspiciousness, which resulted in difficulties in emotional contacts. In this case high levels of neuroticism were observed. During long-term stressful effects the failure of adaptive mechanisms was noted, which often led to disorganisation in educational activities, unpredictability in behaviour (protest, rigidity). This type is not promising for sports.

We judged about the degree of the first-year students' adaptation to the university conditions by the time period of establishing interpersonal

relationships in a group, with students of the same year, in a sports team, by manifestation of organisational skills. The time period served as a system-forming factor characterising the outcome of adaptation.

The success of psycho-physiological adaptation depends on how the student will be able to establish himself as a personality within the group of students, in his interpersonal relations, value orientations, beliefs and internal motivation to study. If the adaptation was successful the student was a quicker learner in terms of self-discipline, self-control, he was able to organise his educational activities, thus contributing to the development of the cognitive activity and intellectual area.

The sprinter (66.8 %) and the mixed (60.0 %) types had the shortest period of adaptation, from 3 to 6 months, i.e. by the time of the first examination session students had set personal relationships in the group, had shown organisational abilities in different events. However, the sprinters in 33.2 % of cases and the mixed type in 40 % of cases during that period of time adapted to the university conditions, but nevertheless set not only positive but also negative (conflict, lack of tolerance) interpersonal relationships, some of them formed informal groups.

The stayers required the period of adaptation from 6 to 12 months, i.e. they got adapted within the first year of studies. Phlegmatic persons were characterised by a low level of commitment to interpersonal relationships, and they preferred a small circle of communication, their low social activity did not contribute to the development of organisational abilities, except for those who were engaged in sports sections. Melancholic persons experienced a necessity for interpersonal relationships, but due to the weakness of the nervous system they did not take the initiative and like phlegmatic persons they sought to limit their social circle, occupying a neutral

position in the group, they displayed sympathy with friends, were drawn to extroverts, but displayed the fear of rejection. The “unsteady type” could not adapt quickly. Adaptation time varied in the range from 12 to 34 months. The person referred to the unsteady type feels uncomfortable, relationships with classmates may be conflicting and ultimately may lead to social and psychological maladjustment. As a result, the student can not reveal his individual capacities, does not fulfill his potential. Hence, the lack of success in studies appears and as a result, the student is dismissed from the university.

The research carried out has shown that the “sprinters” and the “mixed” types adapt equally quickly. Successful adaptation at the end of the semester was observed among the “stayers”. The “unsteady” type is less adaptive, only 68 % out of this group were able to pass the winter session in due time.

The PPA types gain the most significance during the sportsmen training, which has been proved as the result of the long-term experiment on the members of the teams of the Krasnoyarsk Territory and the country in various kinds of sports (Table 3).

During the sportsmen training it is important to consider personality peculiarities and behaviour in competitive conditions. Under prolonged stress effects (long-term training in camps and a lot of starts in different climatic and zone conditions) sportsmen often get a breakdown of adaptive mechanisms. The breakdown of adaptive mechanisms is accompanied by the decrease in efficiency, the disorder of psychomotor functions resulting in the psycho-emotional discomfort and the decrease of sports results.

Timely information about the types of psycho-physiological adaptation allows to adjust the load modes promptly, to optimise

Table 3. Achievements of the sportsmen – participants of the experiment

Family name, first name	Kind of sports	Type of the temperament	Type of psycho-physiological adaptation	Mobility of the nervous processes	Sports qualification	Sports achievements
Chagiyev B.	Greco-Roman style	Sanguine	sprinter	high	Master of sports of the international level	Champion of Europe, of the international tournament named after Poddubny
Sartakov A.	“-“	Phlegmatic	stayer	high	Master of sports of the international level	Champion of the international tournament named after Poddubny
Satiyev B.	freestyle wrestling	Sanguine	sprinter	high	Honoured master of sports	World champion, champion of Europe, Olympic champion 2000, 2004.
Madosyan G.	“-“	Choleric	mixed	good	Master of sports	Silver winner of Yarygin tournament
Fedotkina S.	skating	Sanguine	sprinter	high	Honoured master of sports	Russian champion, silver champion of the Olympiad
Kuts V.	underwater swimming	Sanguine	mixed	high	Master of sports of the international level	Champion of the USSR, RF, winner of international championships, octuple record breaker of the world
Goncharova T.	“-“	Choleric	sprinter	high	Master of sports of the international level	Champion of the RF, the USSR, Europe, sextuple record breaker of the world
Andronov E.	“-“	Sanguine	spinter	high	Master of sports of the international level	Champion of the world, Europe, the RF, octuple record breaker of the world and Europe.

the training process, to select remediation activities, to conduct sessions of mental regulation, thus mobilising functional reserves of the body, and, therefore, allows to avoid the disruption of adaptation and achieve superior results.

From the analysis of the table we can conclude that the type of psycho-physiological adaptation is directly connected with the sports qualification and the results of the sportsman. The

utmost results were achieved by the sportsmen-sprinters of the sanguine type.

Conclusion

We have identified four types of the psycho-physiological adaptation: the “sprinter”, the “stayer”, the “mixed” and the “unsteady” types by using the nonparametric methods of the classification. Using the types of PPA gives the university faculty an opportunity to obtain

objective information on psycho-physiological adaptation of a student and, consequently, to solve not only socially important issues, but also to expand the range of issues which allow to remove the psycho-emotional strain and motivate an individual for maintaining and improving health, increasing the reserve capacity of the body. The obtained experimental data regarding the PPA types are of a great

practical value in sports activities where it is very important to take into account personality peculiarities.

These PPA types belong to one of the prognostic criteria of the psycho-physiological adaptation to extreme conditions, educational, vocational and sports activities, but may also serve as a criterion for sports and professional selection.

References

L.R. Ayrapetyants, *Pedagogicheskiye osnovy planirovaniya i kontrolya sorevnovatelnoy i trenirovochnoy deyatelnosti [Teaching basics of planning and control of competition and training activities]*, State Central Order of Lenin Institute of Physical Culture, M. 1991.

L.R. Ayrapetyants, M.A. Godik, *Sportivnye igry [Sports games]*, Tashkent: Publishing house of Ibn Siny, 1991.

Al-Ghazi Hassan. *Deyatelnost voleibolistov v igrovom protsesse i metodika ee sovershenstvovaniya. [Activities of volleyball players in the game process and methods to improve it]*, Russian State Academy of Physical Culture. M., 1995.

I.P. Andriadi, *Didakticheskiye umeniya trenera i ikh formirovanie. [Didactic coach skills and their formation]*, State Central Order of Lenin Institute of Physical Culture, M., 1983.

G.D. Babushkin, "Vliyaniye sposobnostey, umeniy i otnosheniy trenerov k svoemu delu na effektivnost professionalnoy deyatelnosti. [Impact of abilities, skills, and coach attitude towards work on the efficiency of professional activity]", *Theory and Practice of Physical Culture*, 1985. Is. 1.

A.V. Belyaev, *Kharakteristika sorevnovatelnoy deyatelnosti visokokvalifitsirovannykh volleybolistov. [Characteristics of the competition activity of high-qualified volleyball players]*, M.: State Central Order of Lenin Institute of Physical Culture, 1990.

A.V. Belyaev, *Obucheniye tekhnike igry v voleybol i ee sovershenstvovaniye. [Teaching of the volleyball playing technique and its improvement]*, M.: Human, Olympia, 2007.

A.V. Belyaev, *Pryzhkovaya podgotovka kvalifitsirovannykh voleybolistov v podgotovitelnom periode. Metodicheskaya razrabotka dlya trenerov klubnykh komand I DYUSSH [Jump training of qualified volleyball players during the preparatory period. Methodological development for coaches of club teams and Children and Youth Sports Schools]*, M.: Print Center Ltd, 2004.

S.M. Vaytsekhovskiy, *Kniga trenera [Coach Book]*, M.: Physical Culture and Sport, 1971.

Voleybol: Uchebnik dlya vysshykh uchebnykh zavedeniy fizicheskoy kultury [Volleyball: Textbook for higher educational institutions of physical culture], ed. by A.V. Belyaev, M.V. Savin – M.: Physical culture, Education and Science, 2000.

"Voleybol: Etapy sportivnogo sovershenstvovaniya i vysshego sportivnogo masterstva [Volleyball: Steps in sports perfection and higher levels of sports skills], *Progressive sports training for Children and Youth Sports Schools & Specialized children-youthful school of Olympic reserve*: Ext. State Committee of Russian Federation for Physical Culture and Sport, M.: Sov. Sports, 2004.

I.N. Galaktionov, "Pedagogicheskie sposobnosti trenera [Coach Teaching abilities]", *T&PPC*, 1986. Is. 8.

A.V. Gaskov, *Teoretiko-metodicheskie osnovy upravleniya sorevnovatelnoy i trenirovochnoy deyatel'nostyu kvalifitsirovannykh bokserov. [Theoretical and methodological basics for the management of competition and training activities of the qualified boxers]*, Ulan-Ude: BSU, 1999.

A.Ya. Gomelsky, *Upravlenie komandoy v basketbole [Team management in basketball]*, M.: PC&S, 1985.

V.M. Dyachkov, *Voprosy upravleniya protsessom sovershenstvovaniya sportivnogo masterstva. [Issues of the management process to improve sports skills]*, M.: PC &S, 1972.

N.V. Zhmarev, *Upravlencheskaya i organizatorskaya deyatel'nost trenera [Management and organization of the coach activity]*, Kiev: Zdorov'ya, 1986.

M.Ya. Kim, *Individualnaya igrovaya podgotovka voleybolistok s uchetom osobennosteyresheniya operativnykh zadach. [Individual training of volleyball players in view of peculiarities for solving of the operational objectives]*, Russian State Academy of Physical Culture. – M., 1994.

Yu.N. Kleshchev, *Upravlenie trenirovachnym protsessom v sisteme mnogoletney podgotovki komand vysshykh razryadov po voleybolu. [Training process management in the system of long-term training of volleyball teams of higher categories]*, M.: State Central Order of Lenin Institute of Physical Culture, 1983.

Yu.N. Kleshchev, K.K. Markov, *Voleybol. Kniga trenera. Chast 2 [Volleyball. Coach Book. Part 2]*, Irkutsk: Irkutsk Institute of Railway Engineers, 2000.

N.V. Kuzmina, *Metody issledovaniya pedagogicheskoy deyatel'nosti. [Research methods for the pedagogical activity]*, L., 1970.

H. Lubinska, Faktory, *Obuslavlivayushchie povyshenie effektivnosti deyatel'nosti trenera sportivnoy komandy [Factors responsible for improving the performance of a sports team coach]*, Kiev State Institute of Physical Culture, 1983.

K.K. Markov, "Metodika analiza voleybolnoy tekhniki pri obuchenii i sovershenstvovanii// [Analysis methodology of volleyball techniques in sports training and perfection]", *Proceedings of the Scientific Conference*, Irkutsk State Technical University, 1996.

K.K. Markov, "Strategiya dvigatel'nogo obucheniya v voleybole vysshego urovnya [The strategy of motor education in high-level volleyball training]", *Proceedings of the Scientific Conference*, Irkutsk State Technical University, 1997.

V.N. Maslov, "Pedagogicheskie nabludeniya v protsesse otsenki igrovoy deyatel'nosti voleybolistov [Pedagogical observation in the evaluation process of volleyball players' activity]", *Scientific basics of long-term planning of the training process and preparation of the Olympic reserve*, Dnepropetrovsk: B.i, 1983.

L.P. Matveev, "Problemy vyyavleniya i formirovaniya spetsificheskikh zakonomernostey i printsipov sportivnoy trenirovki [Identification and formation problems of specific laws and principles of sports training]", *Pedagogy, Psychology*, M., 1980.

R.M. Naydiffer, *Psikhologiya sorevnuyushchegosya sportsmena [Psychology of a competing sportsman]*, M.: PC&S, 1979.

M.P. Pimenov, Starichenko L.M. *Upravlenie komandoy v sorevnovaniyakh po voleybolu/ [Team management in volleyball competitions]*, Kiev: KSUPC, 1984.

V.N. Platonov, *Teoriya i metodika sportivnoy trenirovki. [Theory and methods of sports training]*, Kiev: Higher Educational Institution, 1984.

M.V. Prokhorova, *Osobennosti kommunikativnoy deyatel'nosti trenera DUSSCH [Features of the Youth School coach communication]*, State Central Order of Lenin Institute of Physical Culture, M., 1974.

A.Ts. Puni, *Psikhologicheskie voprosy trenirovki i gotovnosti sportsmena k sorevnovaniyu. [Psychological issues of training and sportsman's readiness to compete]*, M.: PC&S, 1969.

A.V. Rodionov, *Psikhologo-pedagogicheskie metody povysheniya effektivnosti operativnoy deyatel'nosti v sporte. [Psychological and pedagogical methods to improve operational effectiveness in sports]*, State Central Order of Lenin Institute of Physical Culture, – M., 1990.

Rukovodstvo trenera po voleibolu/ [Volleyball coach guidance], Translation edited by K.K. Markov, Irkutsk, Irkutsk State Technical University, 1999.

G.I. Savenkov, *Issledovanie psikhologicheskikh osobenostey sostoyaniya trenirovannykh sportsmenov. [Study of psychological features of the trained sportsmen's state]*, State Central Order of Lenin Institute of Physical Culture, M., 1969.

P.V. Simonov, *Vysshaya nervnaya deyatel'nost cheloveka: Motivatsionno-emotsionalnye aspekty. [Human higher nervous activity: Motivational and emotional aspects]* M.: Science, 1975.

Soderzhanie i organizatsiya kompleksnogo kontrolya v podgotovke voleybolistov vysshikh razryadov [The content and organization of integrated control in training of volleyball players of high categories], M.: State Central Order of Lenin Institute of Physical Culture, 1981.

Yu.V. Sysoev, *Stil rukovodstva trenera sportsmenami i sportivnym kolektivom. [Style of coaching sportsmen and sports teams]*, M.: B.i, 1997.

O.P. Topyshev, *Pedagogicheskie aspekty sovershenstvovaniya deyatel'nosti sportsmenov v igrovyykh vidakh sporta [Pedagogical aspects of the improvement of sportsmen activities in playing sports]*, M.: State Central Order of Lenin Institute of Physical Culture, 1989.

V.E. Hapko, *Issledovanie tekhniki igry v voleybols uchetom usloviy izbiratelnykh pedagogicheskikh vozdeystviy [Study of volleyball techniques taking into account conditions of the selective pedagogical effects]*, Kiev, 1977.

O.A. Chernikova, *Emotsii v sporte [Emotions in sports]*, M.: PC&S, 1962.

I. Ahrabi-Fard, "Coaching perceptual preparation in Volleyball" *Seminar on Hitter Development*, Colorado Springs, Feb. 1996.

M. Dudson, "Serving: an antipoden perspective", *The Coach*, 2000, 2.

M. Mohammed, "The role of the coach in youth teams", *The Coach*, 1992, 1.

L. Sawula, *Role of the coach. Operational Manual for FIVB Instructors*, Lausanne: FIVB, 1997.

Психофизиологические типы адаптации в оценке профессионального и спортивного отбора

О.Н. Московченко

*Сибирский федеральный университет
Россия 660041, Красноярск, пр. Свободный, 79*

Предварительными исследованиями были выделены четыре типа психофизиологической адаптации на основе анализа непараметрических алгоритмов автоматической классификации (А.В. Лапко, 1993). Предлагается дополнить ранее выделенные базовые критерии, которые интегрируют нейродинамические профили адаптации, характеризующие индивидуальные особенности центральной нервной системы, ее силу и лабильность, показатели вариабельности сердечного ритма. В качестве дополнительных критериев предлагаются дифференцированные морфофункциональные типы физического развития, а также типологические особенностей личности, определенные по шкале экстра-интроверсия Н. Еysenk (1972). Такой подход правомерно позволит рассматривать типы психофизиологической адаптации в единстве личностных свойств с генетически природными задатками, обеспечивающими индивидуально-психологические особенности личности. Приведена краткая характеристика каждого типа психофизиологической адаптации, что способствует пониманию индивидуальных свойств личности и позволяет спрогнозировать поведенческие реакции.

Проведенные исследования позволили расширить представление о типах психофизиологической адаптации, выделить прогностические критерии, которые позволяют прогнозировать поведение в экстремальных условиях, в условиях учебной, производственной и спортивной деятельности. Типы психофизиологической адаптации правомерно рассматривать при оценке социально-психологической адаптации студентов, а также в качестве критериев профессионального и спортивного отбора.

Ключевые слова: психофизиологические типы адаптации, спринтер, стайер, смешанный тип, ненадежный тип, критерии отбора.
